

Laparoendoscopic single-site hysterectomy in Cuba: a novel technique with new medical instruments

Javier Ernesto Barreras González¹, Yuderkis Castillo Sánchez², Ana Bertha López Milhet², Israel Díaz Ortega², Jorge Gerardo Pereira Fraga², Rafael Torres Peña²

¹Department of Laparoscopic and Endoscopic Surgery, National Center for Minimally Invasive Surgery, Havana, Cuba; ²Department of Laparoscopic and Endoscopic Surgery, National Center for Minimally Invasive Surgery, Havana Medical University, Havana, Cuba

Correspondence to: Javier Ernesto Barreras González, MD, PhD. Department of Laparoscopic and Endoscopic Surgery, National Center for Minimally Invasive Surgery, Párraga Street 315. 10 de octubre. Havana. Cuba. Email: javier.barrera@infomed.sld.cu, javierdeu_3008@yahoo.es.

Abstract: Hysterectomy is a frequent non-obstetrical surgical technique. The main benefits of laparoscopic hysterectomy are less hospital stay and complications, and better surgical results. The aim of this video is to describe a new technique of laparoendoscopic single-site surgery for totally laparoscopic hysterectomy using a laparoscope with a work channel and modified conventional laparoscopic instruments. A 47-year-old female with uterine fibroid underwent laparoendoscopic single-site hysterectomy. The procedure was performed using a laparoscope with a work channel, standard laparoscopic instruments, and modified conventional laparoscopic instruments adapted for these procedures. After general anesthesia, a 10–11-mm trocar was then placed through a vertical umbilical incision. The Clermont Ferrand uterine manipulator was then placed. The procedure was similar to multiport total laparoscopic hysterectomy. Alternative use of bipolar electrocoagulation and scissors was made to gain entry to the cavity by the work channel. Divisions were made in the round ligaments, tube pedicles, and utero-ovarian ligaments by means of bipolar electrocautery. The upper junction of the vesicouterine peritoneal fold was distinguished as a white line and was dissected. Then, dissection of the posterior peritoneum and both uterosacral ligaments was performed. The uterine vessels were dissected safely with bipolar electrocautery. The vagina was opened using a monopolar L-hook over the sealed device. The uterus was removed through the vagina. The vaginal cuff was sutured laparoscopically (only with a needle holder), with one 0 polyglycolic acid suture, using a curved 26–30-mm needle introduced through the vagina and tied extracorporeally through the vagina. A new retraction system for the vagina, designed by the current authors work team, was used. After controlling the bleeding, the laparoscope and the trocar were removed, and the umbilical fascia and subcutaneous tissue were closed. The uterus weight was 300g, operative time was 82 minutes and blood loss was 54 mL. The postoperative hospital stay was less than one day. No morbidity. This technique proposed seems to be a feasible and safe approach and without the disadvantages of loss of triangulation and clashing of instruments, compared with other laparoendoscopic single-site hysterectomy techniques.

Keywords: Laparoscopic hysterectomy; total laparoscopic hysterectomy; laparoendoscopic single-site surgery; minimally invasive surgery; gynecology

Received: 13 December 2020; Accepted: 17 February 2021.

doi: 10.21037/gpm-20-68

View this article at: <http://dx.doi.org/10.21037/gpm-20-68>

Introduction

Hysterectomies are one of the most common surgical procedures worldwide. Following the Caesarean section, it is the second common gynecological surgery in women at

reproductive age (1,2).

For many years, hysterectomy was done via laparotomy or vaginal access (3). To reduce surgical invasion to a minimum, laparoscopic hysterectomy as a less invasive procedure is

developed. Reich (4) in 1989 and Pelosi *et al.* (5) in 1991 reported the first multiport laparoscopic hysterectomy and single-port laparoscopic hysterectomy respectively.

The main benefits of laparoscopic hysterectomy are less hospital stay and complications, and better surgical results (1,6).

Since the introduction of laparoscopic-assisted vaginal hysterectomy (LAVH) at our Hospital in 1994, it became the standard treatment for symptomatic uterine fibroids. When we acquired growing expertise, we began performing totally laparoscopic hysterectomy and laparoendoscopic single-site (LESS) hysterectomy in 1998 and 2010 respectively.

The aim of this video is to describe a new LESS technique for totally laparoscopic hysterectomy (TLH) at the National Center for Minimally Invasive Surgery in Havana, Cuba.

Operative technique

We report a LESS for TLH by means of an operative laparoscope and modified conventional laparoscopic instruments (*Video 1*).

To perform the procedure we utilized a 10–11 mm trocar, an operative laparoscope (0 degrees, 10 mm, with a 5–6 mm working channel), a Clermont Ferrand uterine manipulator, bipolar forceps, prototype of laparoscopic instruments and a new retraction system for vagina.

All procedures performed in studies involving human participants were in accordance with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient for publication of this study and any accompanying images.

Surgical technique

We gave the patient a prophylactic antibiotic while inducing general anesthesia. Vaginal pumping was performed and a urinary catheter was inserted. Patient was placed on the operating table in a modified lithotomy position Surgeon is situated on the left side while first assistant stands between legs. The operative laparoscope was inserted through a vertical umbilical incision and the abdominal and pelvic organs were inspected.

After dilating the cervix (Hegar number 8), intrauterine manipulator is inserted under direct vision.

Surgical technique steps

(I) *Cauterization and transection of the round ligament:* We

used a bipolar electrocoagulation and scissors. The retroperitoneal space is visualized after cutting the round ligament.

- (II) *Cauterization of the tube and the ovarian ligament:* Pedicles and ligaments were coagulated. When adnexectomy was required, we used to retraction the adnexal the same retraction system for the anterior vaginal.
- (III) *Dissection of anterior broad ligament peritoneum:* The anterior broad ligament peritoneum was dissected.
- (IV) *Rejection of the bladder:* After the vesicouterine fold was identified, the bladder was rejected.
- (V) *Dissection of uterosacral ligaments:* Posterior peritoneum was cut and both uterosacral ligaments were dissected.
- (VI) *Dissection and cutting of the uterine vessels:* We used bipolar coagulation for the treatment of uterine vessels.
- (VII) *Opening of vaginal vault and extraction of the uterine specimen:* Vagina was opened by bipolar electrocautery and a monopolar L-hook over the sealed device. After that, the uterus was removed.
- (VIII) *Retraction for vagina (Option 1):* We used a forceps through the vagina to retract the anterior vaginal wall in *the midline*.
- (IX) *Retraction system for vagina (Option 2):* We used a new retraction system to retract anterior vaginal wall.
- (X) *Closure of vaginal vault:* Vaginal vault was sutured laparoscopically using one hand and 0 polyglycolic acid suture.

Retraction system for vagina was released. Suture was tied extracorporeally through the vagina.

Hemostasis was controlled before removing the laparoscope and the trocar.

Postoperative care

After the patient recovered from anesthesia, the urinary catheter was removed. Patient was started with liquid diet 6 hours after surgery and discharged on first postoperative day.

Medical follow-ups were made at 15 days, then at 45 days to visualize the suture of the vaginal vault via speculum.

Results

Patient is a 47-year-old woman with symptomatic uterine fibroid underwent LESS hysterectomy. The uterus weight was 300 g, operation time was 82 minutes and operative blood loss was 54 mL. Length of stay was less than 24 hours.

There was No morbidity.

Comments

Our hospital has the most experience for Minimally Invasive techniques in Cuba. Minimally invasive gynecological team has accomplished over 5,300 endoscopic gynecological procedures, 3,100 TLH and 200 LESS hysterectomies.

LESS-TLH has been proven to be an excellent technique with advantages such as better cosmetic outcomes, less pain, less sepsis and the likelihood of incisional hernias is lower (7-9).

Many studies reported LESS-TLH procedures that utilized multichannel single-incision devices through which the instruments and laparoscope were introduced. Other studies have described the use of many conventional ports inserted by one umbilical incision (7-12).

LESS surgery has a problem with the restraints on instrument manipulation, retraction, and triangulation limits. The collision between the instruments, the instruments with the laparoscope and light cable with the operator's hands occurs because laparoscope and the surgical instrument are positioned in parallel to each other (13). LESS-TLH technique in this study solved those disadvantages.

The operative time was not modified with the suture using one hand and tying it extracorporeally through vagina. This technique has been described for other single-incision laparoscopic surgeries (14). Laparoscopic closure is safer, faster and easier to perform when the technique is standardized which coincides with other reports (15-18).

The vaginal cuff was sutured laparoscopically because laparoscopic approach permits better control of the anterior vaginal wall and the bladder than vaginal approach. We used this close for all laparoscopic hysterectomy (multiport and single port) and several publications show that for vaginal cuff closure, laparoscopic suture is a safe and less time-consuming procedure (compared the vaginal route) and the cuff-related complications are similar in the two groups (15-19).

LESS-TLH proposal is a procedure with the advantages of providing constant vision of the instruments in the operative field and reducing costs (incurred by new devices available today).

Nevertheless, to standardize this technique, some skills must be mastered. First, refers to the need of handling overall optics with the inserted instrument, thus achieving optimal dissection with simultaneous and coordinated movement of the laparoscope laterally and the instruments coaxially. Other elements are related to the work angle of the instruments for dissection regarding the organ. The technique forces the

surgeon to use both hands at the same time.

Researchers reported the good outcomes using a laparoscope with a work channel in LESS procedure (20).

In conclusion, LESS-TLH technique proposal may be generalized because it is a standardized technique as well as a feasible and safe approach that compared other LESS-TLH techniques.

Acknowledgments

The video was awarded the first prize in the Second International Elite Gynecologic Surgery Competition (2020 Masters of Gynecologic Surgery). The authors would like to acknowledge Ricardo Nilo and the department of information and communication technologies of the National Center for Minimally Invasive Surgery in Cuba for helping the authors to complete this video, and Professor Berta Moineiro Plasencia, B.A., for reviewing the English version of this article.

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Gynecology and Pelvic Medicine* for the series "Award-Winning Videos from the Second International Elite Gynecologic Surgery Competition (2020 Masters of Gynecologic Surgery)". The article has undergone external peer review.

Peer Review File: Available at <http://dx.doi.org/10.21037/gpm-20-68>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/gpm-20-68>). The series "Award-Winning Videos from the Second International Elite Gynecologic Surgery Competition (2020 Masters of Gynecologic Surgery)" was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All procedures performed in studies involving human participants were in accordance with the Helsinki Declaration (as revised

in 2013). Written informed consent was obtained from the patient for publication of this study and any accompanying images.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Johnson N, Barlow D, Lethaby A, Tavender E, Curr E, Garry R. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev* 2006;CD003677.
- Nieboer TE, Johnson N, Lethaby A, et al. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev* 2009;3:CD003677.
- Clayton RD. Hysterectomy. *Best Pract Res. Clin Obstet Gynaecol* 2006;20:73-87.
- Reich H, DeCaprio J, McGlynn F. Laparoscopic hysterectomy. *J Gynecol Surg* 1989;5:213-6.
- Pelosi MA, Pelosi MA 3rd. Laparoscopic hysterectomy with bilateral salpingo-oophorectomy using a single umbilical puncture. *N J Med* 1991;88:721.
- Reich H. Total laparoscopic hysterectomy: Indications, techniques and outcomes. *Curr Opin Obstet Gynecol* 2007;19:337.
- Jo EJ, Kim TJ, Lee YY, et al. Laparoendoscopic single-site surgery with hysterectomy in patients with prior cesarean section: Comparison of surgical outcomes with bladder dissection techniques. *J Minim Invasive Gynecol* 2013;20:160.
- Sinha R, Sundaram M, Mahajan C, et al. Single-incision total laparoscopic hysterectomy. *J Minim Access Surg* 2011;7:78.
- Xie W, Cao D, Yang J, et al. Single-Port vs Multiport Laparoscopic Hysterectomy: A Meta-Analysis of Randomized Controlled Trials. *J Minim Invasive Gynecol* 2016;23:1049-56.
- Langebrekke A, Qviqstad E. Total laparoscopic hysterectomy with single-port access without vaginal surgery. *J Minim Invasive Gynecol* 2009;16:609.
- Lee YY, Kim TJ, Kim CJ, et al. Single-port access laparoscopic-assisted vaginal hysterectomy: A novel method with a wound retractor and a glove. *J Minim Invasive Gynecol* 2009;16:450.
- Gou J, Li Z, Liao X, et al. Single-incision laparoscopic hysterectomy using conventional laparoscopic instruments: initial experience with 25 cases. *Arch Gynecol Obstet* 2018;298:921-6.
- Cuschieri A. Single-incision laparoscopic surgery. *J Minim Access Surg* 2011;7:3.
- Thanakumar J, John PH. One-handed knot tying technique in single-incision laparoscopic surgery. *J Minim Access Surg* 2011;7:112.
- Uccella S, Malzoni M, Cromi A, et al. Laparoscopic vs transvaginal cuff closure after total laparoscopic hysterectomy: a randomized trial by the Italian Society of Gynecologic Endoscopy. *Am J Obstet Gynecol* 2018;218:500.e1-500.e13.
- Bastu E, Yasa C, Dural O, et al. Comparison of 2 Methods of Vaginal Cuff Closure at Laparoscopic Hysterectomy and Their Effect on Female Sexual Function and Vaginal Length: A Randomized Clinical Study. *J Minim Invasive Gynecol* 2016;23:986-93.
- Biler A, Kale A, Terzi H, et al. Newly Developed Laparoscopic Needle Holder That Facilitates Knot Tying Makes Vaginal Cuff Suturing Easy in Single-Port Laparoscopic Hysterectomy. *Surg Innov* 2017;24:605-10.
- Terzi H, Biler A, Turkay U, et al. A comparison of novel laparoscopic suturing techniques in single-port surgery. *Minim Invasive Ther Allied Technol* 2019;28:338-43.
- Hwang JH, Lee JK, Lee NW, et al. Vaginal cuff closure: a comparison between the vaginal route and laparoscopic suture in patients undergoing total laparoscopic hysterectomy. *Gynecol Obstet Invest* 2011;71:163-9.
- Khosla A, Ponsky TA. Use of operative laparoscopes in single-port surgery: The forgotten tool. *J Minim Access Surg* 2011;7:116.

doi: 10.21037/gpm-20-68

Cite this article as: Barreras González JE, Sánchez YC, López Milhet AB, Ortega ID, Pereira Fraga JG, Peña RT. Laparoendoscopic single-site hysterectomy in Cuba: a novel technique with new medical instruments. *Gynecol Pelvic Med* 2021.